

Soil and plant testing for profitable fertiliser use

Harm van Rees¹, Sean Mason², Dan Bell³, Therese McBeath⁴, Jackie Ouzman⁴, Rick Llewellyn⁴, Craig Muir⁵, Fiona Tomney⁶ and Ian Richter⁶

¹Cropfacts; ²Agronomy Solutions; ³Landmark; ⁴CSIRO; ⁵Agrivision; ⁶SARDI, Minnipa Agricultural Centre



Key messages

- **Soil P and N status are highly variable across paddocks. Depending on variability, soil sampling intensity should be increased to sample more than one zone in a paddock.**
- **Low production zones tended to have lower soil P and higher soil N levels, suggesting that nutrient inputs might require adjustment to improve profitability.**
- **In 2019 the yield response in the Mallee to P fertiliser was highly variable across the paddock and was closely correlated to soil P status (Colwell with inclusion of PBI interpretation/DGT - proviso, only one paddock had been analysed at the time of writing).**

Why do the trial?

Precision Agriculture for variable rate fertiliser application demands a knowledge of the soil available nutrient variation across a paddock and an understanding of the likely responses to applied nutrients. Soil testing is shifting from surface sampling (0-10 cm) to deep sampling (GRDC Farm Survey 2016) however, farmers and advisers appear to be unsure on how to interpret soil test results to optimise fertiliser returns, especially with variable rate

application of fertiliser. In 2016, it was estimated 15% of paddocks were regularly tested, as opposed to 40% in 2008 (GRDC pers. comm.)

How was it done?

Landmark, independent consultants and Farming Systems Groups including EPARF are partners in the project to raise understanding and awareness around issues dealing with nitrogen (N) and phosphorus (P) responses with variable rate fertiliser application, inclusive of undertaking intensive soil testing in different production zones across paddocks. APAL laboratories are undertaking the soil and plant analysis. CSIRO are analysing yield maps, performing the statistical analysis of yields achieved on the P and N rate strips, and reviewing the economic implications of 'informed' P rate applications based on soil testing.

Paddock trials 2019

Over 300 paddock-based trials were established in 2019 in SA and Victoria from close to 700 sampling zones. Production zones in paddocks were defined either by using historical yield maps or the farmers' long-term knowledge of the paddock. For two production zones in each paddock, a 1 ha soil sampling area was selected – the two zones were located in-line with the sowing direction. Sampling intensity for each 1 ha soil sampling area were 36 topsoil samples for available P (0-10 cm: Colwell, DGT, PBI) and six deep cores (0-10, 10-30, 30-60, 60-90 cm) for available N (NO₃ and NH₄) with the samples combined for

each depth to generate one soil test value. Chloride was included in the analysis to determine whether sub-soil salinity inhibited yield.

In 150 of the 333 paddocks sampled in 2019, farmers sowed P rate strips across the paddock, ensuring the strips crossed the 1 ha soil sampling grids. Available soil P status and likely fertiliser P response rates were calculated from Colwell and DGT in association with PBI. The rates of P applied were informed by the soil test result and most sown strips included a 0 rate, farmer rate and double the farmer rate of applied P in situations predicted to be P responsive, and the inclusion of a half farmer rate for situations predicted to be non-P responsive. The P strips received the same N as applied by the farmer for the rest of the paddock. Tissue samples to check on tissue P status and possible nutrient deficiencies along with dry matter estimates were collected from each fertiliser rate strip between growth stage (GS) 16 and 32.

Fertiliser N response

In 2019 a number of paddocks also had different top-dressed N strips applied to generate N rate trials in paddocks where soil N variability was high, these were applied at the same time as the farmers applied in-crop urea. As with the P scenario, N trials had rates of N applied as informed by the starting soil N profile and most sown strips included a 0 rate, farmer rate and double the farmer rate of applied N in responsive situations, and the inclusion of half farmer rate for non-responsive situations.

Harvest and statistical analysis

Yield monitor data were used to calculate the yield for each P and N fertiliser rate strip. The yield achieved for each fertiliser rate strip within each 1 ha soil sampling area was used to correlate crop yield to soil P and N status. Harvest strip data within each of the two soil sampling zones was analysed for significant difference using a moving average t-test (Lawes and Bramley 2012) enabling the evaluation of nutrient treatment responses between zones and within zones. A partial gross margin analysis will be undertaken to calculate the change in income achieved from the different fertiliser rate strips.

What happened?

Soil N and P status 2019

A brief snapshot of the nutrient status across the Southern region revealed high variability of both N and P between the two production

zones in each sampled paddock. There were opportunities identified within each agroecological zone for the establishment of both N and P trials. As an overall summary the N status was generally good and supported at least the production of a 2 t/ha wheat crop (Figure 1), without factoring in immobilisation/mineralisation. In general, the N status was higher for the low production areas which indicates N build up due to lower N removal caused by a soil constraint or low yields in seasons prior.

At a paddock level, P deficiency is driven by the ability of different soils to fix/absorb P sources as estimated from the PBI (Phosphorus Buffer Index). Quite often low production zones were associated with low extractable P, high PBI and relatively high soil N due to less utilisation of N sources and subsequent removal (Figure 2). In these circumstances

simple replacement fertiliser strategies are unbalanced and are creating a wider gap between yield production zones.

Improved gross margins from more efficient fertiliser applications are expected if different production zones are assessed for the ability of the soil to provide the crop with adequate nutrients.

Paddock trials 2019

An example of the experimentation is presented for a paddock sown to wheat (Scepter - sowing date 15/5/2019) in the Victorian Mallee.

Soil P results

Soil P results for Colwell and DGT P, and PBI (Phosphorus Buffering Index, see Burkitt *et al.* 2002) are detailed in Table 1. In Zone 1 both soil tests predicted marginal P while in Zone 2 the DGT P soil test predicted deficient soil P. PBI was relatively high in Zone 2.




Figure 1. Overall soil mineral N status across the Southern region for allocated high production and low production zones within paddocks before the 2019 sowing season. Using the rule of thumb of 40 kg N/ha required for 1 t/ha grain, the low production zone supported 2.0 t/ha compared with 2.5 t/ha for the high production zone. Error bars represents standard error across all sampling sites in each zone.





Figure 2. Overall soil P status across the southern region for allocated high and low production zones within each paddock as assessed by Colwell P (left) and DGT (right) together with PBI for each zone. Critical Colwell P was determined by the relationship generated in Moody, 2007. Critical DGT value for wheat is 64 ug/L (95% CI = 53-78 ug/L). Error bars represents standard error across all sampling sites in each zone.

Table 1. Mallee paddock - P test result pre-sowing 2019 (Colwell, DGT and PBI) for Zone 1 and 2.

P Test	Zone 1	Zone 2
Colwell P (mg/kg)	22	30
DGT P (ug/L)	42	12
PBI	64	135




Figure 3. Strip yield (t/ha) for two rates of fertiliser P applied from paddock boundary to boundary, crossing two soil sampling areas. Solid black circles and squares indicate the yield achieved within the soil sampling areas for Zone 1 and 2.

Table 2. Yield response to four rates of fertiliser P applied at sowing in two zones.

Rate (P kg/ha)	Average yield (t/ha) within production zone		Average yield (t/ha) entire strip
	Zone 1	Zone 2	
0	2.51	1.76 ^a	2.16
4.4	2.60	1.76 ^a	2.24
8.8	2.59	1.67 ^a	2.01
17.6	2.32	2.34 ^b	2.22
Sign. difference	ns	P<0.05	

Zone 2: ^a and ^b denote significant difference

P rate trial

Four rates of P (as MESZ) were applied at sowing with double seeder width strips across the paddock through each zone (fertiliser applied at 0, 4.4, 8.8 and 17.6 kg P/ha) (all strips had urea at 45 kg/ha, 20.7 kg N/ha, applied at sowing). Urea at 75 kg/ha (34.5 kg N/ha) was top-dressed on the trial area on 28/6/2019.

Harvest yield map data was used to analyse the yield differences between P treatments in each of the two soil sampling areas (1 ha areas located in two distinct production zones in line of sowing). Statistical analysis was based on the t-test for comparing two strips (example Figure 3).

A significant difference in yield gain was found only in Zone 2 for the high rate of P applied (17.6 kg P/ha) (Table 2).

What does this mean?

Soil nutrient status is highly variable across paddocks and these initial results indicate that we need to sample more than one soil type/production zone in a paddock.

Indicative results indicate that intensive soil sampling of different production zones provides significant benefit in terms of P application (results from the N rate application strips had not been analysed at the time of writing).

This research project is ongoing until 2022, so further information and results will be available for paddocks monitored on Eyre Peninsula during the season.

References

- Burkitt LL, Moody PW, Gourley CJP and Hannah MC 2002. A simple phosphorus buffering index for Australian soils. Aust. J. Soil Research 40, 497-513
- Lawes RA and Bramley RGV 2012. A simple method for the analysis of on-farm strip trials. Agronomy Journal 104: 371-377
- Moody, PW 2007. Interpretation of a single-point P buffering index for adjusting critical levels of the Colwell soil P test. Australian Journal Soil Research 45: 55-62

Acknowledgements

The research undertaken as part of this project is made possible by the significant contributions of growers through both trial cooperation and the support of the GRDC (GRDC project 9176604), the authors thank the GRDC for their continued support. We also acknowledge the farmers who are in this project for sowing the strip trials, applying variable rate P and N, and supplying yield monitor data at harvest. Through EPARF, the EP the growers and families involved in the project are Wes Matthews (Koongawa), Terry and Thomas Schmucker (Koongawa), Peter Kuhlmann (Mudamuckla), Andrew and Tim Polkinghorne (Lock) and the Minnipa Agricultural Centre staff.



AGRONOMY SOLUTIONS

INDEPENDENT RESEARCH • QUALITY RESULTS



Eyre Peninsula Agricultural Research Foundation Inc.



SARDI

